

CONVERGENCE ANALYSIS FOR A STABILIZED LINEAR SEMI-IMPLICIT NUMERICAL SCHEME FOR THE NONLOCAL CAHN–HILLIARD EQUATION

XIAO LI, ZHONGHUA QIAO, AND CHENG WANG

ABSTRACT. In this paper, we provide a detailed convergence analysis for a first order stabilized linear semi-implicit numerical scheme for the nonlocal Cahn–Hilliard equation, which follows from consistency and stability estimates for the numerical error function. Due to the complicated form of the nonlinear term, we adopt the discrete H^{-1} norm for the error function to establish the convergence result. In addition, the energy stability obtained by Du et al., [J. Comput. Phys. 363 (2018), pp. 39–54] requires an assumption on the uniform ℓ^∞ bound of the numerical solution, and such a bound is figured out in this paper by conducting the higher order consistency analysis. Taking the view that the numerical solution is indeed the exact solution with a perturbation, the error function is ℓ^∞ bounded uniformly under a loose constraint of the time step size, which then leads to the uniform maximum-norm bound of the numerical solution.

1. INTRODUCTION

In this paper, our primary purpose is to develop a detailed convergence analysis of a stabilized linear semi-implicit numerical scheme for the nonlocal Cahn–Hilliard (NCH) equation taking the form [4, 11, 24]

$$(1) \quad u_t = \Delta(u^3 - u + \varepsilon^2 \mathcal{L}u), \quad (\mathbf{x}, t) \in \Omega \times (0, T],$$

where $u = u(\mathbf{x}, t)$ is the unknown function subject to the periodic boundary condition. Here, $\Omega = \prod_{i=1}^d (-X_i, X_i)$ is a rectangular domain in \mathbb{R}^d , $T > 0$ is the terminal time, $\varepsilon > 0$ is an interfacial parameter, and \mathcal{L} is a nonlocal linear operator defined by

$$(2) \quad \mathcal{L} : v(\mathbf{x}) \mapsto \int_{\Omega} J(\mathbf{x} - \mathbf{y})(v(\mathbf{x}) - v(\mathbf{y})) \, d\mathbf{y},$$

where J is a kernel function satisfying following conditions [11]:

- (a) $J(\mathbf{x}) \geq 0$ for any $\mathbf{x} \in \Omega$;
- (b) J is even, i.e., $J(\mathbf{x}) = J(-\mathbf{x})$ for any $\mathbf{x} \in \mathbb{R}^d$;

Received by the editor November 13, 2018, and, in revised form, January 5, 2020, and March 16, 2020.

2010 *Mathematics Subject Classification.* Primary 35Q99, 65M12, 65M15, 65M70.

Key words and phrases. Nonlocal Cahn–Hilliard equation, Stabilized linear scheme, convergence analysis, higher order consistency expansion.

The first author’s work was partially supported by NSFC grant 11801024.

The second author’s work was partially supported by the Hong Kong Research Council GRF grants 15325816 and 15300417.

The third author’s work was partially supported by NSF grant NSF DMS-1418689.

The second author is the corresponding author.

- (c) J is Ω -periodic;
- (d) $\frac{1}{2} \int_{\Omega} J(\mathbf{x}) |\mathbf{x}|^2 d\mathbf{x} = 1$,

where condition (d) means that the kernel has a finite second order moment in Ω . In fact, J could be taken as a radial function over the domain $\Omega = \prod_{i=1}^d (-X_i, X_i)$, with exponential decay at the boundary, such as $J(\mathbf{x}) = \alpha \exp\left(-\frac{|\mathbf{x}|^2}{\sigma^2}\right)$ (for a small σ), and a periodic extension is made to \mathbb{R}^d , so that both (b) and (c) are satisfied. The NCH equation (1) can be viewed as the H^{-1} gradient flow with respect to the free energy functional

$$(3) \quad E(u) = \int_{\Omega} F(u(\mathbf{x})) d\mathbf{x} + \frac{\varepsilon^2}{2} (\mathcal{L}u, u)_{L^2},$$

with $F(u) = \frac{1}{4}(u^2 - 1)^2$, or equivalently, by using condition (b),

$$(4) \quad E(u) = \int_{\Omega} \left(F(u(\mathbf{x})) + \frac{\varepsilon^2}{4} \int_{\Omega} J(\mathbf{x} - \mathbf{y}) (u(\mathbf{x}) - u(\mathbf{y}))^2 d\mathbf{y} \right) d\mathbf{x}.$$

The second term in (4) usually represents the interaction energy, describing the long-range interactions between atoms at different sites, and the kernel J measures the strength of interactions. Using the Taylor formula, the periodicity of u , and conditions (b)–(d) of J , one can show that

$$\frac{\varepsilon^2}{4} \int_{\Omega} J(\mathbf{x} - \mathbf{y}) (u(\mathbf{x}) - u(\mathbf{y}))^2 d\mathbf{y} \approx \frac{\varepsilon^2}{2} |\nabla u(\mathbf{x})|^2,$$

which suggests that the classic Cahn–Hilliard equation [9]

$$(5) \quad u_t = \Delta(u^3 - u - \varepsilon^2 \Delta u),$$

corresponding to the local energy functional

$$(6) \quad E_{\text{local}}(u) = \int_{\Omega} \left(F(u(\mathbf{x})) + \frac{\varepsilon^2}{2} |\nabla u(\mathbf{x})|^2 \right) d\mathbf{x},$$

is an approximation of the NCH equation (1) under the assumption that the interaction exists only in a very short range.

If J is further integrable, then $J * 1 = \int_{\Omega} J(\mathbf{x}) d\mathbf{x} > 0$ is a positive constant and

$$(7) \quad \mathcal{L}v = (J * 1)v - J * v,$$

where

$$(J * v)(\mathbf{x}) = \int_{\Omega} J(\mathbf{x} - \mathbf{y}) v(\mathbf{y}) d\mathbf{y} = \int_{\Omega} J(\mathbf{y}) v(\mathbf{x} - \mathbf{y}) d\mathbf{y}$$

is exactly the periodic convolution [24]. In this case, the NCH equation (1) can be written as

$$u_t = \nabla \cdot (a(u) \nabla u) - \varepsilon^2 \Delta J * u,$$

where $a(u) = 3u^2 - 1 + \varepsilon^2 J * 1$ is referred to as diffusive mobility. If

$$(8) \quad \gamma_0 := \varepsilon^2 J * 1 - 1 > 0,$$

which gives $a(u) > 0$, then equation (1) becomes diffusive and the solution becomes regular in time; otherwise, the solution may exhibit some singular behavior. Throughout this paper, we always assume that the kernel J is integrable with condition (8) held.

As one of the typical systems of the phase field models, the classic Cahn–Hilliard equation (5) has been successfully used to model phase transitions occurring in

mixtures of small molecules and some other interface problems involving mass-conserved order parameters. Recently, the NCH equation (1) has attracted increasing attention and been used in various fields ranging from materials science to finance and image processing. For instance, in materials science, the NCH equation and other related equations arise as mesoscopic models of interacting particle systems [2, 25] and are taken to model phase transitions [18]; in the dynamic density functional theory [1, 2], the interaction kernel is the two-particle direct correlation function and the solution represents the mesoscopic particle density. In the theoretical level, the well-posedness of the NCH equations equipped with a Neumann or Dirichlet boundary condition were investigated by Bates and Han [6, 7] by assuming the integrability of the kernel. Du et al. [10] developed a general framework of nonlocal diffusion problems, and a number of examples ranging from continuum mechanics to graph theory were showed to be special cases of the proposed framework. For more details on theoretical investigations, see [4, 8, 18–21].

There have been several works on numerical analysis for nonlocal models. For a class of nonlocal diffusion models with various boundary conditions, finite difference and finite element approximations were addressed in [35, 36, 44]. For the nonlocal Allen–Cahn equation, the L^2 gradient flow with respect to (4), Bates et al. [5] developed an L^∞ stable and convergent finite difference scheme by treating the nonlinear and nonlocal terms explicitly, and Du et al. [13] analyzed the spectral-Galerkin approximations. In addition, the maximum principle preserving property has been established for the exponential time differencing (ETD) schemes in a more recent work [12]. For the NCH equation, an important fact is that the exact solution decreases the energy in time due to the energetic variational structure of the underlying model, so it is highly desirable to develop numerical algorithms inheriting this property of energy stability at the discrete level. Energy stability has been widely investigated for numerical schemes of a family of classic PDE-based phase field models, such as convex splitting schemes [17, 29, 42], stabilized schemes [33, 43], and so on. The application of similar analysis for nonlocal phase field models is still full of challenges due to the lack of the higher order diffusion term. Guan et al. [22–24] constructed convex splitting schemes for the NCH equation and the nonlocal Allen–Cahn equation by treating the nonlinear term implicitly and putting the nonlocal term into the explicit part. The proposed scheme allows one to evaluate the nonlocal term explicitly only once at each time step, but iterations are inevitable due to the nonlinearity of the scheme.

In order to avoid the nonlinear iteration, in a recent work [11] a linear semi-implicit scheme has been developed by using the stabilizing approach. The linear nonlocal term is set in the implicit level and solved efficiently in the frequency space by using the fast Fourier transform (FFT) technique due to the linearity of the resulted fully discrete system. The first order stabilized linear semi-implicit (SSI1) scheme given in [11] reads

$$(9) \quad \frac{u^{n+1} - u^n}{\Delta t} = \Delta_N [(u^n)^3 - u^n + A(u^{n+1} - u^n) + \varepsilon^2 \mathcal{L}_N u^{n+1}],$$

and the energy stability has been proved, that is, $E_N(u^{n+1}) \leq E_N(u^n)$ if the stabilizing constant A satisfies

$$(10) \quad A \geq \frac{1}{2} \|u^{n+1}\|_\infty^2 + \|u^n\|_\infty^2 - \frac{1}{2}.$$

Here, E_N , Δ_N , and \mathcal{L}_N are the spatially discretized forms of the operators E , Δ , and \mathcal{L} , respectively, and their precise definitions will be given in the next section. Notice that the infinity-norms of the numerical solutions at time steps t_n and t_{n+1} have been involved on the right-hand side of (10). However, such a lower bound for constant A has not been justified.

We aim to justify the lower bound of A in this paper. A direct analysis provided in [26–28] for the local Cahn–Hilliard model could hardly be extended to the case of nonlocal models due to the lack of higher order diffusion terms. Instead, we view the numerical solution as a perturbation of the exact solution to (1), perform a local-in-time convergence analysis, and obtain the ℓ^∞ bound of the numerical solution via the convergence result. All the analysis will be specified in the 2-D case; similar results can be obtained for the 1-D and 3-D cases without any extra essential difficulties.

The outline of the paper is as follows. Some notation and lemmas for the spectral collocation method for the spatial discretization are summarized in Section 2. The convergence analysis, as well as the ℓ^∞ bound of the numerical solutions, of the first order stabilized linear semi-implicit scheme (9) is presented in Section 3. Finally, some concluding remarks are given in Section 4.

2. SPECTRAL COLLOCATION METHOD FOR THE SPATIAL DISCRETIZATION

In this section, we summarize some notation and lemmas introduced in [11] for the spectral collocation approximations of some spatial operators in the two-dimensional space with $\Omega = (-X, X) \times (-Y, Y)$.

Let N_x and N_y be two even numbers. The $N_x \times N_y$ mesh Ω_h of the domain Ω is a set of nodes (x_i, y_j) with $x_i = -X + ih_x$, $y_j = -Y + jh_y$, $1 \leq i \leq N_x$, $1 \leq j \leq N_y$, where $h_x = 2X/N_x$ and $h_y = 2Y/N_y$ are the uniform mesh sizes in each direction. Let $h = \max\{h_x, h_y\}$. We define the index sets

$$S_h = \{(i, j) \in \mathbb{Z}^2 \mid 1 \leq i \leq N_x, 1 \leq j \leq N_y\},$$

$$\widehat{S}_h = \left\{ (k, l) \in \mathbb{Z}^2 \mid -\frac{N_x}{2} + 1 \leq k \leq \frac{N_x}{2}, -\frac{N_y}{2} + 1 \leq l \leq \frac{N_y}{2} \right\}.$$

All of the periodic grid functions defined on Ω_h are denoted by \mathcal{M}_h , that is,

$$\mathcal{M}_h = \{f : \Omega_h \rightarrow \mathbb{R} \mid f_{i+mN_x, j+nN_y} = f_{ij} \text{ for any } (i, j) \in S_h \text{ and } (m, n) \in \mathbb{Z}^2\}.$$

For any $f, g \in \mathcal{M}_h$ and $\mathbf{f} = (f^1, f^2)^T$, $\mathbf{g} = (g^1, g^2)^T \in \mathcal{M}_h \times \mathcal{M}_h$, the discrete L^2 inner product $\langle \cdot, \cdot \rangle$, discrete L^2 norm $\|\cdot\|_2$, and discrete L^∞ norm $\|\cdot\|_\infty$ are, respectively, defined by

$$\begin{aligned} \langle f, g \rangle &= h_x h_y \sum_{(i,j) \in S_h} f_{ij} g_{ij}, & \langle \mathbf{f}, \mathbf{g} \rangle &= h_x h_y \sum_{(i,j) \in S_h} (f_{ij}^1 g_{ij}^1 + f_{ij}^2 g_{ij}^2), \\ \|f\|_2 &= \sqrt{\langle f, f \rangle}, & \|\mathbf{f}\|_2 &= \sqrt{\langle \mathbf{f}, \mathbf{f} \rangle}, \\ \|f\|_\infty &= \max_{(i,j) \in S_h} |f_{ij}|, & \|\mathbf{f}\|_\infty &= \max_{(i,j) \in S_h} \sqrt{|f_{ij}^1|^2 + |f_{ij}^2|^2}. \end{aligned}$$

For any $f \in \mathcal{M}_h$, we call $\bar{f} := \frac{1}{4XY} \langle f, 1 \rangle$ the mean value of f . In particular, denote by \mathcal{M}_h^0 all the grid functions in \mathcal{M}_h with mean zero, i.e.,

$$\mathcal{M}_h^0 = \{f \in \mathcal{M}_h \mid \langle f, 1 \rangle = 0\}.$$

2.1. Discrete gradient, divergence, and Laplace operators. For a function $f \in \mathcal{M}_h$, the 2-D discrete Fourier transform $\hat{f} = Pf$ is defined componentwise [32, 37] by

$$(11) \quad \hat{f}_{kl} = \sum_{(i,j) \in S_h} f_{ij} \exp\left(-i \frac{k\pi}{X} x_i\right) \exp\left(-i \frac{l\pi}{Y} y_j\right), \quad (k, l) \in \hat{S}_h.$$

The function f can be reconstructed via the corresponding inverse transform $f = P^{-1}\hat{f}$ with components given by

$$(12) \quad f_{ij} = \frac{1}{N_x N_y} \sum_{(k,l) \in \hat{S}_h} \hat{f}_{kl} \exp\left(i \frac{k\pi}{X} x_i\right) \exp\left(i \frac{l\pi}{Y} y_j\right), \quad (i, j) \in S_h.$$

Let $\widehat{\mathcal{M}}_h = \{Pf \mid f \in \mathcal{M}_h\}$ and define the operators \widehat{D}_x and \widehat{D}_y on $\widehat{\mathcal{M}}_h$ as

$$(\widehat{D}_x \hat{f})_{kl} = \left(\frac{k\pi i}{X}\right) \hat{f}_{kl}, \quad (\widehat{D}_y \hat{f})_{kl} = \left(\frac{l\pi i}{Y}\right) \hat{f}_{kl}, \quad (k, l) \in \hat{S}_h.$$

Then the Fourier spectral approximations to the first and second partial derivatives can be represented as

$$D_x = P^{-1} \widehat{D}_x P, \quad D_y = P^{-1} \widehat{D}_y P, \quad D_x^2 = P^{-1} \widehat{D}_x^2 P, \quad D_y^2 = P^{-1} \widehat{D}_y^2 P.$$

For any $f \in \mathcal{M}_h$ and $\mathbf{f} = (f^1, f^2)^T \in \mathcal{M}_h \times \mathcal{M}_h$, the discrete gradient, divergence, and Laplace operators are given, respectively, by

$$\nabla_N f = \begin{pmatrix} D_x f \\ D_y f \end{pmatrix}, \quad \nabla_N \cdot \mathbf{f} = D_x f^1 + D_y f^2, \quad \Delta_N f = D_x^2 f + D_y^2 f.$$

It is easy to prove the following results.

Lemma 1. (i) For any $f, g \in \mathcal{M}_h$ and $\mathbf{g} \in \mathcal{M}_h \times \mathcal{M}_h$, we have the summation by parts formulas

$$\langle f, \nabla_N \cdot \mathbf{g} \rangle = -\langle \nabla_N f, \mathbf{g} \rangle, \quad \langle f, \Delta_N g \rangle = -\langle \nabla_N f, \nabla_N g \rangle = \langle \Delta_N f, g \rangle.$$

(ii) The inversion of $-\Delta_N$ exists on \mathcal{M}_h^0 and $(-\Delta_N)^{-1}$ is self-adjoint and positive definite.

Lemma 1(ii) tells us that $(-\Delta_N)^{-1}f$ is well-defined for any $f \in \mathcal{M}_h^0$. Then we can define the discrete H^{-1} norm $\|\cdot\|_{-1,N}$ by

$$(13) \quad \|f\|_{-1,N} = \sqrt{\langle f, (-\Delta_N)^{-1}f \rangle} \quad \forall f \in \mathcal{M}_h^0.$$

2.2. Discrete convolution and nonlocal operators. To define the discrete convolutions, we consider the kernel function set

$$\mathcal{K}_h = \{\psi : \Omega_{h,0} \rightarrow \mathbb{R} \mid \psi_{i+mN_x, j+nN_y} = \psi_{ij} \text{ for any } (i, j) \in S_h \text{ and } (m, n) \in \mathbb{Z}^2\},$$

where $\Omega_{h,0} = \{(ih_x, jh_y) \mid (i, j) \in S_h\}$ is the mesh on the domain $(0, 2X) \times (0, 2Y)$. A discrete transform and its inversion of a function $\psi \in \mathcal{K}_h$ could be defined similarly via (11) and (12) by replacing x_i and y_j by ih_x and jh_y , respectively. Actually, \mathcal{K}_h is equivalent to \mathcal{M}_h due to the periodicity of their elements, and we consider the functions from \mathcal{K}_h as the kernels just for convenience of notation.

For any $\psi \in \mathcal{K}_h$ and $f \in \mathcal{M}_h$, the discrete convolution $\psi \circledast f \in \mathcal{M}_h$ is defined componentwise by

$$(\psi \circledast f)_{ij} = h_x h_y \sum_{(m,n) \in S_h} \psi_{i-m, j-n} f_{mn}, \quad (i, j) \in S_h.$$

Especially, by setting $f \equiv 1$ on Ω_h , we have

$$\psi \circledast 1 = h_x h_y \sum_{(m,n) \in S_h} \psi_{mn}.$$

The following preliminary estimate is needed in the convergence analysis.

Lemma 2. Suppose $J \in C_{\text{per}}^1(\Omega)$ and define its grid restriction by $J_{ij} := J(x_i, y_j)$. Then for any $f, g \in \mathcal{M}_h$, we have

$$(14) \quad |\langle J \circledast f, \Delta_N g \rangle| \leq \alpha \|f\|_2^2 + \frac{C}{\alpha} \|\nabla_N g\|_2^2,$$

for any $\alpha > 0$, where C is a positive constant that depends on J but is independent of h_x and h_y .

Proof. An application of summation by parts and the Cauchy–Schwarz inequality shows that

$$(15) \quad |\langle J \circledast f, \Delta_N g \rangle| = |\langle \nabla_N(J \circledast f), \nabla_N g \rangle| \leq \|\nabla_N(J \circledast f)\|_2 \cdot \|\nabla_N g\|_2.$$

An application of the definitions of discrete gradient and convolution gives us

$$(16) \quad (\nabla_N(J \circledast f))_{ij} = h_x h_y \sum_{(m,n) \in S_h} (\nabla_N J)_{i-m, j-n} f_{mn}, \quad (i, j) \in S_h.$$

Then,

$$\begin{aligned} \|\nabla_N(J \circledast f)\|_2^2 &= h_x h_y \sum_{(i,j) \in S_h} \left(h_x h_y \sum_{(m,n) \in S_h} (\nabla_N J)_{i-m, j-n} f_{mn} \right)^2 \\ &\leq |\Omega| \|\nabla_N J\|_\infty^2 \left(h_x h_y \sum_{(m,n) \in S_h} f_{mn} \right)^2 \\ &\leq |\Omega|^2 \|\nabla_N J\|_\infty^2 \|f\|_2^2. \end{aligned}$$

The smoothness of J implies the bound $\|\nabla_N J\|_\infty \leq C_J$. Then, we arrive at

$$(17) \quad |\langle J \circledast f, \Delta_N g \rangle| \leq C_J |\Omega| \|f\|_2 \|\nabla_N g\|_2 \leq \alpha \|f\|_2^2 + \frac{C}{\alpha} \|\nabla_N g\|_2^2,$$

for any $\alpha > 0$, where $C = \frac{1}{4} C_J^2 |\Omega|^2$. \square

Remark 1. We make a technical assumption $J \in C_{\text{per}}^1(\Omega)$ to facilitate the analysis. On the other hand, a singular J may also be of certain scientific interest in many relevant physical models. However, a direct application of the above analysis is not available for a singular J , since a pointwise bound of $\nabla_N J$ is not available in (16) any more. A nonstandard extension to the case with a singular J will be considered in our future works.

Given an integrable kernel J satisfying assumptions (a)–(d), we can define the discrete version of the nonlocal operator \mathcal{L} by

$$(18) \quad \mathcal{L}_N f = (J \circledast 1)f - J \circledast f \quad \forall f \in \mathcal{M}_h.$$

It is easy to check that \mathcal{L}_N commutes with Δ_N and is self-adjoint and positive semidefinite. Finally, the discrete version of the energy (3) is defined as

$$(19) \quad E_N(v) = \langle F(v), 1 \rangle + \frac{\varepsilon^2}{2} \langle \mathcal{L}_N v, v \rangle, \quad v \in \mathcal{M}_h.$$

2.3. Fourier projection of the exact solution. The existence and uniqueness of a smooth periodic solution to the Integro-Partial Differential Equation (IPDE) (1) with smooth periodic initial data may be established by using techniques developed by Bates and Han in [6, 7]. In this article, we denote this IPDE solution by U . Motivated by these results, one obtains

$$(20) \quad \|U\|_{L^\infty(0,T;L^\infty)} + \|\nabla U\|_{L^\infty(0,T;L^\infty)} < C,$$

for any $T > 0$.

Define $U_N(\cdot, t) := \mathcal{P}_N U(\cdot, t)$, the (spatial) Fourier projection of the exact solution into \mathcal{B}^N , the space of trigonometric polynomials of degree up to N . The following projection approximation is standard: if $U \in L^\infty(0, T; H_{\text{per}}^\ell)$, for some $\ell \in \mathbb{N}$,

$$(21) \quad \|U_N - U\|_{L^\infty(0,T;H^k)} \leq Ch^{\ell-k} \|U\|_{L^\infty(0,T;H^\ell)}, \quad h = \max\{h_x, h_y\} \quad \forall 0 \leq k \leq \ell.$$

By U_N^m , U^m we denote $U_N(\cdot, t_m)$ and $U(\cdot, t_m)$, respectively, with $t_m = m\Delta t$. It is clear that $\int_\Omega U_N(\cdot, t_m) d\mathbf{x} = \int_\Omega U(\cdot, t_m) d\mathbf{x}$, for any $m \in \mathbb{N}$, due to the fact that U_N is the Fourier projection of U , and thus,

$$(22) \quad \int_\Omega U_N(\cdot, t_m) d\mathbf{x} = \int_\Omega U(\cdot, t_m) d\mathbf{x} = \int_\Omega U(\cdot, t_{m-1}) d\mathbf{x} = \int_\Omega U_N(\cdot, t_{m-1}) d\mathbf{x} \quad \forall m \in \mathbb{N},$$

in which the second step is based on the fact that the exact solution U is mass conservative at the continuous level. On the other hand, the solution of the numerical scheme (9) is also mass conservative at the discrete level:

$$(23) \quad \overline{u^m} = \overline{u^{m-1}} \quad \forall m \in \mathbb{N}.$$

Meanwhile, we denote by u_N^m the values of U_N at discrete grid points at time instant t_m , i.e., $u_N^m := \mathcal{P}_h U_N(\cdot, t_m)$. Since $U_N \in \mathcal{B}^N$, it always holds that

$$\int_\Omega U_N(\cdot, t_m) d\mathbf{x} = h_x h_y \sum_{(i,j) \in S_h} U_N(x_i, y_j, t_m) = h_x h_y \sum_{(i,j) \in S_h} (u_N^m)_{i,j},$$

so the mass conservative property is available at the discrete level: $\overline{u_N^m} = \overline{u_N^{m-1}}$. As indicated before, we use the mass conservative projection for the initial data: $u^0 = u_N^0 = \mathcal{P}_h U_N(\cdot, t = 0)$, that is,

$$(24) \quad u_{i,j}^0 := U_N(x_i, y_j, t = 0).$$

The error grid function is defined as

$$(25) \quad e^m := u_N^m - u^m \quad \forall m \geq 0.$$

Therefore, it follows that

$$(26) \quad \overline{e^m} = 0, \quad \text{since } \overline{u_N^m} = \overline{u_N^0} = \overline{u^0} = \overline{u^m} \quad \forall m \geq 0,$$

so that the discrete norm $\|\cdot\|_{-1,N}$ is well defined for the error grid function. We also notice that the Fourier projection of the exact solution has to be taken at the initial time step as in (24), instead of a pointwise interpolation of the exact initial data, to assure the zero-mean property of the numerical error grid function at a

discrete level. In addition, we have (with $N_k := \lfloor \frac{T}{\Delta t} \rfloor$ denoting the integer part of $\frac{T}{\Delta t}$)

$$(27) \quad \max_{1 \leq k \leq N_k} \|u_N^k\|_\infty + \max_{1 \leq k \leq N_k} \|\nabla_N u_N^k\|_\infty < C.$$

3. CONVERGENCE ANALYSIS AND ENERGY STABILITY ANALYSIS

We begin this section by stating the main result on the convergence analysis of the stabilized linear scheme (9). The detailed proof is given in the following two subsections, including the higher order consistency analysis and the convergence analysis. The energy stability of (9) is then obtained under some new assumptions on A , instead of (10) given in [11]. With an initial data with sufficient regularity, we could assume that the exact solution has regularity of class \mathcal{R} :

$$(28) \quad U \in \mathcal{R} := H^4(0, T; C_{\text{per}}^0) \cap H^3(0, T; C_{\text{per}}^2) \cap L^\infty(0, T; C_{\text{per}}^{m+2}), \quad m \geq 3.$$

Theorem 1. *Given the periodic initial data $U(x, y, t = 0) \in C_{\text{per}}^{m+2}(\Omega)$, suppose the unique periodic solution for the IPDE (1), given by $U(x, y, t)$ on $\Omega \times (0, T]$ for some $T < \infty$, is of regularity class \mathcal{R} . In addition, the following assumption is made for the constant A :*

$$(29) \quad A \geq \frac{18M_0^4}{\gamma_0}, \quad \text{with } M_0 = 1 + \max_{1 \leq k \leq N_k} \|u_N^k\|_\infty.$$

Then, provided Δt and h are sufficiently small, under linear refinement path constraint $\Delta t \leq Ch$, with C any fixed constant, we have

$$(30) \quad \|e^n\|_{-1, N} + \left(\gamma_0 \Delta t \sum_{k=1}^n \|e^k\|_2^2 \right)^{\frac{1}{2}} \leq C(\Delta t + h^m),$$

for all positive integers n such that $n\Delta t \leq T$, where $C > 0$ is independent of h and Δt .

3.1. Higher order consistency analysis. By consistency, the Fourier projection solution U_N solves the discrete equation with an $O(\Delta t + h^m)$ accuracy:

$$(31) \quad \frac{U_N^{n+1} - U_N^n}{\Delta t} = \Delta_N \left((U_N^n)^3 - U_N^n + A(U_N^{n+1} - U_N^n) + \varepsilon^2 \mathcal{L}_N U_N^{n+1} \right) + \tau_0^{n+1},$$

where the local truncation error τ_0^{n+1} satisfies

$$(32) \quad \|\tau_0^{n+1}\|_{-1, N} \leq C(\Delta t + h^m).$$

In addition, the discrete zero-mean property of τ_0^{n+1} is observed, which will be useful in later analysis:

$$(33) \quad \overline{\tau_0^{n+1}} = 0,$$

since

$$\overline{U_N^{n+1}} = \overline{U_N^n}, \quad \int_{\Omega} \Delta_N \left((U_N^n)^3 - U_N^n + A(U_N^{n+1} - U_N^n) + \varepsilon^2 \mathcal{L}_N U_N^{n+1} \right) d\mathbf{x} = 0.$$

Notice that the first identity is derived in (22), while the second one comes from the periodic boundary condition. However, this local truncation error will not be enough to recover the $\|\cdot\|_\infty$ bound of the numerical solution due to the first order

accuracy in time. To remedy this, we have to construct supplementary fields, $U_{\Delta t}^1$, $U_{\Delta t}^2$, and denote

$$(34) \quad \hat{U} = U_N + \Delta t \mathcal{P}_N U_{\Delta t}^1 + \Delta t^2 \mathcal{P}_N U_{\Delta t}^2.$$

We also notice that both $U_{\Delta t}^1$, $U_{\Delta t}^2$ are (spatially) continuous functions, and their construction will be outlined later. Moreover, a higher $O(\Delta t^3 + h^m)$ consistency has to be satisfied with the given numerical scheme (9). The constructed fields $U_{\Delta t}^1$, $U_{\Delta t}^2$, which will be obtained using a perturbation expansion, will depend solely on the exact solution U .

In other words, we introduce a higher order approximate expansion of the exact solution, since a first order temporal consistency estimate (32) is not able to control the discrete ℓ^∞ norm of the numerical solution. Instead of substituting the exact solution into the numerical scheme, a careful construction of an approximate profile is performed by adding $O(\Delta t)$ and $O(\Delta t^2)$ correction terms to the exact solution to satisfy an $O(\Delta t^3)$ truncation error. In turn, we estimate the numerical error function between the constructed profile and the numerical solution, instead of a direct comparison between the numerical solution and exact solution. Such a higher order consistency enables us to derive a higher order convergence estimate in the $\|\cdot\|_{-1,N}$ norm, which in turn leads to a desired $\|\cdot\|_\infty$ bound of the numerical solution, via an application of inverse inequality. This approach has been reported for a wide class of nonlinear PDEs; see the related works for the incompressible fluid equation [15, 16, 30, 31, 38–40], various gradient equations [3, 22, 24], the porous medium equation based on the energetic variational approach [14], nonlinear wave equation [41], etc.

We begin with an application of the temporal discretization in the numerical scheme (9) for the Fourier projection solution U_N :

$$(35) \quad \frac{U_N^{n+1} - U_N^n}{\Delta t} = \Delta \left((U_N^n)^3 - U_N^n + A(U_N^{n+1} - U_N^n) + \varepsilon^2 \mathcal{L} U_N^{n+1} \right) + \Delta t \mathbf{g}^{(1)} + O(\Delta t^2) + O(h^m),$$

which comes from the Taylor expansion in time. In more detail, the function $\mathbf{g}^{(1)}$ is smooth enough and only depends on the higher order derivatives of U_N . In particular, by making use of similar arguments as in the derivation of (33), we conclude that

$$(36) \quad \int_{\Omega} (U_N^{n+1} - U_N^n) d\mathbf{x} = 0, \quad \int_{\Omega} \Delta \left((U_N^n)^3 - U_N^n + A(U_N^{n+1} - U_N^n) + \varepsilon^2 \mathcal{L} U_N^{n+1} \right) d\mathbf{x} = 0.$$

This in turn indicates that

$$(37) \quad \int_{\Omega} \mathbf{g}^{(1)} d\mathbf{x} = 0.$$

The first order temporal correction function $U_{\Delta t}^1$ is given by the solution of the following linear differential equation:

$$(38) \quad \partial_t U_{\Delta t}^1 = \Delta \left(3U_N^2 U_{\Delta t}^1 - U_{\Delta t}^1 + \varepsilon^2 \mathcal{L} U_{\Delta t}^1 \right) - \mathbf{g}^{(1)},$$

$$(39) \quad U_{\Delta t}^1(\cdot, t = 0) \equiv 0,$$

with the periodic boundary condition. In fact, (38) is a linear parabolic PDE, with a sufficiently regular coefficient function $3U_N^2$ (regularity dependent on $U \in \mathcal{R}$). The existence and uniqueness of its solution could be derived by making use of a standard

Galerkin procedure and Sobolev estimates, following the classical techniques for a time-dependent parabolic equation [34]. Such a solution depends solely on the profile U_N and is regular enough. Similar to (35), an application of the temporal discretization to $U_{\Delta t}^1$ indicates that

$$(40) \quad \frac{(U_{\Delta t}^1)^{n+1} - (U_{\Delta t}^1)^n}{\Delta t} = \Delta \left(3(U_N^n)^2 (U_{\Delta t}^1)^n - (U_{\Delta t}^1)^n + A((U_{\Delta t}^1)^{n+1} - (U_{\Delta t}^1)^n) + \varepsilon^2 \mathcal{L}(U_{\Delta t}^1)^{n+1} \right) - (\mathbf{g}^{(1)})^n + O(\Delta t).$$

In turn, we denote $\hat{U}^{(1)} = U_N + \Delta t \mathcal{P}_N U_{\Delta t}^1$. A combination of (35) and a Fourier projection of (40) results in the following higher order consistency estimate:

$$(41) \quad \frac{(\hat{U}^{(1)})^{n+1} - (\hat{U}^{(1)})^n}{\Delta t} = \Delta \left(((\hat{U}^{(1)})^n)^3 - (\hat{U}^{(1)})^n + A((\hat{U}^{(1)})^{n+1} - (\hat{U}^{(1)})^n) + \varepsilon^2 \mathcal{L}(\hat{U}^{(1)})^{n+1} \right) + \Delta t^2 \mathbf{g}^{(2)} + O(\Delta t^3) + O(h^m),$$

in which we have made use of the following estimate:

$$(42) \quad \begin{aligned} (\hat{U}^{(1)})^3 &= (U_N + \Delta t \mathcal{P}_N U_{\Delta t}^1)^3 = U_N^3 + 3\Delta t U_N^2 \mathcal{P}_N U_{\Delta t}^1 + O(\Delta t^2) \\ &= U_N^3 + 3\Delta t \mathcal{P}_N (U_N^2 \mathcal{P}_N U_{\Delta t}^1) + O(\Delta t^2) + O(h^m). \end{aligned}$$

Again, $\mathbf{g}^{(2)}$ is smooth enough and only dependent on the higher order derivatives of U_N .

In addition, we observe that the constructed profile $U_{\Delta t}^1$ has zero-mean at the continuous level, based on equations (38)–(39), combined with the fact (37):

$$(43) \quad \int_{\Omega} \partial_t U_{\Delta t}^1 d\mathbf{x} = \int_{\Omega} \Delta \left(3U_N^2 U_{\Delta t}^1 - U_{\Delta t}^1 + \varepsilon^2 \mathcal{L} U_{\Delta t}^1 \right) d\mathbf{x} - \int_{\Omega} \mathbf{g}^{(1)} d\mathbf{x} = 0,$$

so that

$$(44) \quad \int_{\Omega} U_{\Delta t}^1(\cdot, t) d\mathbf{x} = \int_{\Omega} U_{\Delta t}^1(\cdot, t=0) d\mathbf{x} = 0 \quad \forall t > 0.$$

In turn, its projection also has a zero-mean:

$$(45) \quad \int_{\Omega} \mathcal{P}_N U_{\Delta t}^1(\cdot, t) d\mathbf{x} = 0 \quad \forall t > 0.$$

Therefore, we conclude that $\hat{U}^{(1)}$ has the same average as U_N at the continuous level:

$$(46) \quad \int_{\Omega} \hat{U}^{(1)}(\cdot, t) d\mathbf{x} = \int_{\Omega} U_N(\cdot, t) d\mathbf{x}, \quad \forall t > 0.$$

Since U_N is mass conservative at the continuous level, as indicated by (22), we arrive at a similar property for $\hat{U}^{(1)}$:

$$(47) \quad \int_{\Omega} (\hat{U}^{(1)})^{n+1} d\mathbf{x} = \int_{\Omega} (\hat{U}^{(1)})^n d\mathbf{x} \quad \forall n \in \mathbb{N}.$$

As a consequence, by making use of similar arguments as in (36)–(37), we see that $\mathbf{g}^{(2)}$ has zero-mean at the continuous level:

$$(48) \quad \int_{\Omega} \mathbf{g}^{(2)} d\mathbf{x} = 0.$$

The second order temporal correction function $U_{\Delta t}^2$ could be constructed in a similar manner, and it turns out to be the solution of the following linear differential equation:

$$(49) \quad \partial_t U_{\Delta t}^2 = \Delta \left(3U_N^2 U_{\Delta t}^2 - U_{\Delta t}^2 + \varepsilon^2 \mathcal{L} U_{\Delta t}^2 \right) - \mathbf{g}^{(2)},$$

$$(50) \quad U_{\Delta t}^2(\cdot, t = 0) \equiv 0,$$

with the periodic boundary condition. Similarly, (49) is a linear parabolic PDE, with a sufficiently regular coefficient function $3U_N^2$, and its unique solution depends solely on the profile U_N and is smooth enough. An application of the temporal discretization to $U_{\Delta t}^2$ gives

$$(51) \quad \frac{(U_{\Delta t}^2)^{n+1} - (U_{\Delta t}^2)^n}{\Delta t} = \Delta \left(3(U_N^n)^2 (U_{\Delta t}^2)^n - (U_{\Delta t}^2)^n + A((U_{\Delta t}^2)^{n+1} - (U_{\Delta t}^2)^n) \right. \\ \left. + \varepsilon^2 \mathcal{L} (U_{\Delta t}^2)^{n+1} \right) - (\mathbf{g}^{(2)})^n + O(\Delta t).$$

Notice that $\hat{U} = \hat{U}^{(1)} + \Delta t^2 \mathcal{P}_N U_{\Delta t}^2$. In turn, a combination of (41) and (51) leads to the desired third order consistency estimate in time:

$$(52) \quad \frac{\hat{U}^{n+1} - \hat{U}^n}{\Delta t} = \Delta \left((\hat{U}^n)^3 - \hat{U}^n + A(\hat{U}^{n+1} - \hat{U}^n) + \varepsilon^2 \mathcal{L} \hat{U}^{n+1} \right) + O(\Delta t^3) + O(h^m),$$

in which we have made use of the following estimate:

$$(53) \quad \hat{U}^3 = (\hat{U}^{(1)} + \Delta t^2 \mathcal{P}_N U_{\Delta t}^2)^3 = (\hat{U}^{(1)})^3 + 3\Delta t^2 U_N^2 \mathcal{P}_N U_{\Delta t}^2 + O(\Delta t^3) \\ = (\hat{U}^{(1)})^3 + 3\Delta t^2 \mathcal{P}_N (U_N^2 \mathcal{P}_N U_{\Delta t}^2) + O(\Delta t^3) + O(h^m).$$

Similar to the analyses in (43)–(47), we are able to prove that \hat{U} has the same average as U_N at the continuous level:

$$(54) \quad \int_{\Omega} \partial_t U_{\Delta t}^2 d\mathbf{x} = \int_{\Omega} \Delta \left(3U_N^2 U_{\Delta t}^2 - U_{\Delta t}^2 + \varepsilon^2 \mathcal{L} U_{\Delta t}^2 \right) d\mathbf{x} - \int_{\Omega} \mathbf{g}^{(2)} d\mathbf{x} = 0,$$

so that

$$(55) \quad \int_{\Omega} U_{\Delta t}^2(\cdot, t) d\mathbf{x} = \int_{\Omega} U_{\Delta t}^2(\cdot, t = 0) d\mathbf{x} = 0 \quad \forall t > 0,$$

$$(56) \quad \int_{\Omega} \mathcal{P}_N U_{\Delta t}^2(\cdot, t) d\mathbf{x} = 0 \quad \forall t > 0,$$

$$(57) \quad \int_{\Omega} \hat{U}(\cdot, t) d\mathbf{x} = \int_{\Omega} \hat{U}^{(1)}(\cdot, t) d\mathbf{x} = \int_{\Omega} U_N(\cdot, t) d\mathbf{x} \quad \forall t > 0,$$

$$(58) \quad \int_{\Omega} \hat{U}^{n+1} d\mathbf{x} = \int_{\Omega} \hat{U}^n d\mathbf{x} \quad \forall n \in \mathbb{N}.$$

Finally, with an application of Fourier pseudo-spectral approximation in space, we obtain the $O(\Delta t^3 + h^m)$ truncation error estimate for the constructed solution \hat{U} :

$$(59) \quad \frac{\hat{U}^{n+1} - \hat{U}^n}{\Delta t} = \Delta_N \left((\hat{U}^n)^3 - \hat{U}^n + A(\hat{U}^{n+1} - \hat{U}^n) + \varepsilon^2 \mathcal{L}_N \hat{U}^{n+1} \right) + \tau_2^{n+1},$$

$$(60) \quad \text{with } \|\tau_2^{n+1}\|_{-1,N} \leq C(\Delta t^3 + h^m).$$

We notice that τ_2^{n+1} has zero-mean at a discrete level, $\overline{\tau_2^{n+1}} = 0$, for any $n \in \mathbb{N}$, based on the estimate (58), combined with the fact that $\hat{U}^k \in \mathcal{B}^N$.

As stated earlier, the purpose of the higher order expansion (34) is to obtain an ℓ^∞ bound of the error function via its $\|\cdot\|_{-1,N}$ norm in higher order accuracy by utilizing an inverse inequality in spatial discretization, which will be shown below. A detailed analysis shows that

$$(61) \quad \|\hat{U} - U_N\|_\infty \leq C\Delta t,$$

since $\|U_{\Delta t}^1\|_\infty, \|U_{\Delta t}^2\|_\infty \leq C$. In particular, the following bound becomes available:

$$(62) \quad \|\hat{U} - U_N\|_\infty \leq C\Delta t \leq \frac{1}{2},$$

provided that Δt is sufficiently small, so that

$$(63) \quad \|\hat{U}\|_\infty \leq \|U_N\|_\infty + \|\hat{U} - U_N\|_\infty \leq \|U_N\|_\infty + \frac{1}{2}.$$

3.2. Convergence analysis in the $\ell^\infty(0, T; H_h^{-1}) \cap \ell^2(0, T; \ell^2)$ norm. Instead of a direct comparison between the numerical solution and the Fourier projection U_N of the exact solution, we estimate the error between the numerical solution and the constructed solution to obtain a higher order convergence in the $\|\cdot\|_{-1,N}$ norm. In turn, the following error function is introduced:

$$(64) \quad \hat{e}^k := \hat{U}^k - u^k.$$

In particular, the established consistency estimate (57) indicates that

$$(65) \quad \overline{\hat{U}^k} = \frac{1}{|\Omega|} \int_\Omega \hat{U}^k d\mathbf{x} = \frac{1}{|\Omega|} \int_\Omega U_N^k d\mathbf{x} \quad \forall k \in \mathbb{N},$$

in which the first step is based on the fact that $\hat{U}^k \in \mathcal{B}^N$. Its combination with (26) results in the discrete zero-mean property of the numerical error function \hat{e}^k :

$$(66) \quad \overline{\hat{e}^k} = 0, \quad \text{since } \overline{\hat{U}^k} = \overline{U_N^k} = \overline{U_N^0} = \overline{u^0} = \overline{u^k} \quad \forall k \geq 0.$$

In turn, the discrete $\|\cdot\|_{-1,N}$ norm of this error function is well defined.

Subtracting (9) from (59) yields

$$(67) \quad \frac{\hat{e}^{n+1} - \hat{e}^n}{\Delta t} = \Delta_N \left((\hat{U}^n)^3 - (u^n)^3 - \hat{e}^n + A(\hat{e}^{n+1} - \hat{e}^n) + \varepsilon^2 \mathcal{L}_N \hat{e}^{n+1} \right) + \tau_2^{n+1}.$$

To carry out the nonlinear error estimate, we make a $\|\cdot\|_\infty$ assumption for the numerical error function at the previous time step t_n :

$$(68) \quad \|\hat{e}^n\|_\infty \leq \frac{1}{2}.$$

In turn, the $\|\cdot\|_\infty$ bound for the numerical solution at t^n becomes available:

$$(69) \quad \|u^n\|_\infty = \|\hat{U}^n - \hat{e}^n\|_\infty \leq \|\hat{U}^n\|_\infty + \|\hat{e}^n\|_\infty \leq \|U_N^n\|_\infty + \frac{1}{2} + \frac{1}{2} \leq M_0,$$

in which the estimate (62) for $\|\hat{U}^n\|_\infty$ has been recalled in the third step. The a priori assumption (68) will be recovered in the convergence estimate at the next time step, as will be demonstrated later.

Since $\overline{e^k} = 0$ for any $k \geq 0$, $(-\Delta_N)^{-1}\hat{e}^k$ has been well-defined. Taking a discrete inner product with (67) by $2(-\Delta_N)^{-1}\hat{e}^{n+1}$ leads to

$$(70) \quad \begin{aligned} & \|\hat{e}^{n+1}\|_{-1,N}^2 - \|\hat{e}^n\|_{-1,N}^2 + \|\hat{e}^{n+1} - \hat{e}^n\|_{-1,N}^2 + 2A\Delta t \langle \hat{e}^{n+1} - \hat{e}^n, \hat{e}^{n+1} \rangle \\ &= -2\Delta t \langle (\hat{U}^n)^3 - (u^n)^3, \hat{e}^{n+1} \rangle + 2\Delta t \langle \hat{e}^n, \hat{e}^{n+1} \rangle - 2\varepsilon^2 \Delta t \langle \mathcal{L}_N \hat{e}^{n+1}, \hat{e}^{n+1} \rangle \\ & \quad + 2\Delta t \langle (-\Delta_N)^{-1} \hat{e}^{n+1}, \tau_2^{n+1} \rangle, \end{aligned}$$

in which summation by parts formulas have been repeatedly applied. For the left-hand side term associated with the artificial regularization, the following identity is valid:

$$(71) \quad 2A \langle \hat{e}^{n+1} - \hat{e}^n, \hat{e}^{n+1} \rangle = A(\|\hat{e}^{n+1}\|_2^2 - \|\hat{e}^n\|_2^2 + \|\hat{e}^{n+1} - \hat{e}^n\|_2^2).$$

The right-hand side term associated with the truncation error could be bounded in a straightforward way:

$$(72) \quad 2 \langle (-\Delta_N)^{-1} \hat{e}^{n+1}, \tau_2^{n+1} \rangle \leq 2 \|\hat{e}^{n+1}\|_{-1,N} \cdot \|\tau_2^{n+1}\|_{-1,N} \leq \|\hat{e}^{n+1}\|_{-1,N}^2 + \|\tau_2^{n+1}\|_{-1,N}^2.$$

For the second linear term on the right-hand side, a direct application of Cauchy inequality gives

$$(73) \quad 2 \langle \hat{e}^n, \hat{e}^{n+1} \rangle \leq \|\hat{e}^n\|_2^2 + \|\hat{e}^{n+1}\|_2^2.$$

For the nonlocal linear term on the right-hand side, we begin with a rewritten form:

$$(74) \quad \begin{aligned} -2\varepsilon^2 \langle \mathcal{L}_N \hat{e}^{n+1}, \hat{e}^{n+1} \rangle &= -2\varepsilon^2 \langle (J \circledast 1) \hat{e}^{n+1} - J \circledast \hat{e}^{n+1}, \hat{e}^{n+1} \rangle \\ &= -2\varepsilon^2 (J \circledast 1) \|\hat{e}^{n+1}\|_2^2 + 2\varepsilon^2 \langle J \circledast \hat{e}^{n+1}, \hat{e}^{n+1} \rangle. \end{aligned}$$

Meanwhile, for the term $2\varepsilon^2 \langle J \circledast \hat{e}^{n+1}, \hat{e}^{n+1} \rangle$, we apply (14) in Lemma 2 and obtain

$$(75) \quad \begin{aligned} 2\varepsilon^2 \langle J \circledast \hat{e}^{n+1}, \hat{e}^{n+1} \rangle &= -2\varepsilon^2 \langle J \circledast \hat{e}^{n+1}, \Delta_N ((-\Delta_N)^{-1} \hat{e}^{n+1}) \rangle \\ &\leq \frac{\gamma_0}{2} \|\hat{e}^{n+1}\|_2^2 + \frac{C_3}{\gamma_0} \|\nabla_N (-\Delta_N)^{-1} \hat{e}^{n+1}\|_2^2 \\ &\leq \frac{\gamma_0}{2} \|\hat{e}^{n+1}\|_2^2 + \frac{C_3}{\gamma_0} \|\hat{e}^{n+1}\|_{-1,N}^2, \end{aligned}$$

with C_3 only depending on C_2 and ε . Subsequently, a combination of (74) and (75) yields

$$(76) \quad -2\varepsilon^2 \langle \mathcal{L}_N \hat{e}^{n+1}, \hat{e}^{n+1} \rangle \leq -2\varepsilon^2 (J \circledast 1) \|\hat{e}^{n+1}\|_2^2 + \frac{\gamma_0}{2} \|\hat{e}^{n+1}\|_2^2 + \frac{C_3}{\gamma_0} \|\hat{e}^{n+1}\|_{-1,N}^2.$$

For the nonlinear inner product on the right-hand side of (70), we begin with a rewritten form:

$$(77) \quad -2 \langle (\hat{U}^n)^3 - (u^n)^3, \hat{e}^{n+1} \rangle = -2 \langle (\hat{U}^n)^3 - (u^n)^3, \hat{e}^n \rangle - 2 \langle (\hat{U}^n)^3 - (u^n)^3, \hat{e}^{n+1} - \hat{e}^n \rangle.$$

Because of the nonlinear expansion

$$(78) \quad (\hat{U}^n)^3 - (u^n)^3 = ((\hat{U}^n)^2 + \hat{U}^n u^n + (u^n)^2) \hat{e}^n,$$

we see that the first term on the right-hand side of (77) is always nonpositive:

$$(79) \quad -2 \langle (\hat{U}^n)^3 - (u^n)^3, \hat{e}^n \rangle = -2 \langle ((\hat{U}^n)^2 + \hat{U}^n u^n + (u^n)^2), (\hat{e}^n)^2 \rangle \leq 0.$$

The other term on the right-hand side of (77) could be represented as

$$(80) \quad -2 \langle (\hat{U}^n)^3 - (u^n)^3, \hat{e}^{n+1} - \hat{e}^n \rangle = -2 \langle ((\hat{U}^n)^2 + \hat{U}^n u^n + (u^n)^2) \hat{e}^n, \hat{e}^{n+1} - \hat{e}^n \rangle.$$

On the other hand, the $\|\cdot\|_\infty$ estimate (63) for \hat{U} and the a priori bound (69) have implied that

$$(81) \quad \|\hat{U}^n\|_\infty \leq M_0, \quad \|u^n\|_\infty \leq M_0.$$

These facts yield the following estimate:

$$(82) \quad \|(\hat{U}^n)^2 + \hat{U}^n u^n + (u^n)^2\|_\infty \leq 3M_0^2.$$

In turn, we obtain the following inequality:

$$(83) \quad \begin{aligned} -2\langle (\hat{U}^n)^3 - (u^n)^3, \hat{e}^{n+1} - \hat{e}^n \rangle &\leq 2\|(\hat{U}^n)^2 + \hat{U}^n u^n + (u^n)^2\|_\infty \cdot \|\hat{e}^n\|_2 \cdot \|\hat{e}^{n+1} - \hat{e}^n\|_2 \\ &\leq 6M_0^2 \|\hat{e}^n\|_2 \cdot \|\hat{e}^{n+1} - \hat{e}^n\|_2 \\ &\leq \frac{\gamma_0}{2} \|\hat{e}^n\|_2^2 + \frac{18M_0^4}{\gamma_0} \|\hat{e}^{n+1} - \hat{e}^n\|_2^2. \end{aligned}$$

As a consequence, a substitution of (79) and (83) into (77) gives

$$(84) \quad -2\langle (\hat{U}^n)^3 - (u^n)^3, \hat{e}^{n+1} \rangle \leq \frac{\gamma_0}{2} \|\hat{e}^n\|_2^2 + \frac{18M_0^4}{\gamma_0} \|\hat{e}^{n+1} - \hat{e}^n\|_2^2.$$

Therefore, a substitution of (71)–(73), (76), and (84) into (70) results in

$$(85) \quad \begin{aligned} &\|\hat{e}^{n+1}\|_{-1,N}^2 - \|\hat{e}^n\|_{-1,N}^2 + A\Delta t(\|\hat{e}^{n+1}\|_2^2 - \|\hat{e}^n\|_2^2) \\ &\quad + \left(A - \frac{18M_0^4}{\gamma_0}\right)\Delta t\|\hat{e}^{n+1} - \hat{e}^n\|_2^2 + \left(2\varepsilon^2(J \otimes 1) - 1 - \frac{\gamma_0}{2}\right)\Delta t\|\hat{e}^{n+1}\|_2^2 \\ &\leq \left(1 + \frac{\gamma_0}{2}\right)\Delta t\|\hat{e}^n\|_2^2 + \left(1 + \frac{C_3}{\gamma_0}\right)\Delta t\|\hat{e}^{n+1}\|_{-1,N}^2 + \Delta t\|\tau_2^{n+1}\|_{-1,N}^2. \end{aligned}$$

Under the constraint (29) for the regularization parameter A , and making use of the diffusivity condition (8), we get

$$(86) \quad \begin{aligned} &\|\hat{e}^{n+1}\|_{-1,N}^2 - \|\hat{e}^n\|_{-1,N}^2 + A\Delta t(\|\hat{e}^{n+1}\|_2^2 - \|\hat{e}^n\|_2^2) + \left(1 + \frac{3\gamma_0}{2}\right)\Delta t\|\hat{e}^{n+1}\|_2^2 \\ &\leq \left(1 + \frac{\gamma_0}{2}\right)\Delta t\|\hat{e}^n\|_2^2 + \left(1 + \frac{C_3}{\gamma_0}\right)\Delta t\|\hat{e}^{n+1}\|_{-1,N}^2 + \Delta t\|\tau_2^{n+1}\|_{-1,N}^2. \end{aligned}$$

Subsequently, an application of discrete Gronwall inequality results in the desired convergence estimate:

$$(87) \quad \|\hat{e}^{n+1}\|_{-1,N} + \left(\gamma_0 \Delta t \sum_{k=1}^{n+1} \|\hat{e}^k\|_2^2\right)^{1/2} \leq C^*(\Delta t^3 + h^m),$$

due to the fact that $\|\tau_2^k\|_{-1,N} \leq C(\Delta t^3 + h^m)$, for $k \leq n+1$.

Moreover, we have to recover the a priori assumption (68) at time instant t_{n+1} , so that the analysis could be carried out in the induction style. An application of an inverse inequality to the convergence estimate (87) implies that

$$(88) \quad \begin{aligned} \|\hat{e}^{n+1}\|_\infty &\leq \frac{C\|\hat{e}^{n+1}\|_{-1,N}}{h^2} \leq \frac{CC^*(\Delta t^3 + h^m)}{h^2} \leq \frac{C'C^*(h^3 + h^m)}{h^2} \\ &\leq \frac{C_4 C^* h^3}{h^2} = C_4 C^* h \leq \frac{1}{2}, \quad \text{provided that } h \leq \frac{1}{2C_4 C^*}, \end{aligned}$$

in which we have used the linear refinement path constraint $\Delta t \leq Ch$, as well as the fact that $m \geq 3$. This completes the error estimate for \hat{e} , the numerical error between the numerical solution ϕ , and the constructed approximation solution \hat{U} .

Finally, the error estimate (30) is a direct consequence of the identity

$$(89) \quad e^k = \hat{e}^k - \Delta t U_{\Delta t}^1 - \Delta t^2 U_{\Delta t}^2,$$

which comes from the construction (34), as well as the fact that

$$(90) \quad \|(U_{\Delta t}^1)^k\|_2 \leq C, \quad \|(U_{\Delta t}^2)^k\|_2 \leq C, \quad \text{for any } k \geq 0.$$

The proof for Theorem 1 is completed.

3.3. Theoretical justification of the energy stability. It has been proved in [11] that the energy stability for the numerical scheme (9) is valid under condition (10). In addition, the convergence analysis reveals that the $\|\cdot\|_\infty$ bound (81) for the numerical solution is available as long as another constraint (29) for A is valid, so that the convergence analysis could pass through. The following corollary provides a theoretical justification of the energy stability.

Corollary 1. *Under the assumptions of Theorem 1, the energy stability, namely, $E_N(u^{n+1}) \leq E_N(u^n)$, is valid, under the following constraint for the regularization parameter A :*

$$(91) \quad A \geq \max\left\{\frac{18M_0^4}{\gamma_0}, \frac{3}{2}M_0^2 - \frac{1}{2}\right\}, \quad \text{with } M_0 = 1 + \max_{1 \leq k \leq N_k} \|u_N^k\|_\infty.$$

4. CONCLUDING REMARKS

In this work, we present detailed error estimates for a first order stabilized semi-implicit numerical scheme for the nonlocal Cahn–Hilliard equation, where the Fourier pseudospectral method is used for the spatial discretization. We consider the discrete H^{-1} norm for the error function to establish the convergence result, which avoids the complicated analysis on the nonlinear term. In order to bound the error function in the ℓ^∞ norm, we combine the standard technique for the convergence analysis with a higher order consistency estimate to ensure the convergence order high enough to use the inverse inequality. As a result of the ℓ^∞ boundness of the error function, we derive the uniform ℓ^∞ bound of the numerical solution, and then, the energy stability of the numerical scheme, obtained in [11], is improved by requiring a new assumption on the stabilizer.

It is worth mentioning that we use the higher order consistency analysis to pick up only the temporal truncated error since the spatial spectral accuracy $O(h^m)$ is sufficient as long as m is large enough. However, if one considers the lower order spatial approximations, for instance, the finite difference and finite element methods, the truncated error is usually of the order two and the higher order consistency estimate is also necessary to pick up the spatial truncated error; see, e.g., [22, 24] and the references therein.

REFERENCES

- [1] A. Archer and R. Evans, *Dynamical density functional theory and its application to spinodal decomposition*, J. Chem. Phys. **121** (2004), 4246–4254.
- [2] A. J. Archer and M. Rauscher, *Dynamical density functional theory for interacting Brownian particles: stochastic or deterministic?*, J. Phys. A **37** (2004), no. 40, 9325–9333, DOI 10.1088/0305-4470/37/40/001. MR2095422
- [3] A. Baskaran, J. S. Lowengrub, C. Wang, and S. M. Wise, *Convergence analysis of a second order convex splitting scheme for the modified phase field crystal equation*, SIAM J. Numer. Anal. **51** (2013), no. 5, 2851–2873, DOI 10.1137/120880677. MR3118257

- [4] P. W. Bates, *On Some Nonlocal Evolution Equations Arising in Materials Science*, Nonlinear Dynamics and Evolution Equations, Fields Inst. Commun., vol. 48, Amer. Math. Soc., Providence, RI, 2006, pp. 13–52. MR2223347
- [5] P. W. Bates, S. Brown, and J. Han, *Numerical analysis for a nonlocal Allen-Cahn equation*, Int. J. Numer. Anal. Model. **6** (2009), no. 1, 33–49. MR2574896
- [6] P. W. Bates and J. Han, *The Dirichlet boundary problem for a nonlocal Cahn-Hilliard equation*, J. Math. Anal. Appl. **311** (2005), no. 1, 289–312, DOI 10.1016/j.jmaa.2005.02.041. MR2165478
- [7] P. W. Bates and J. Han, *The Neumann boundary problem for a nonlocal Cahn-Hilliard equation*, J. Differential Equations **212** (2005), no. 2, 235–277, DOI 10.1016/j.jde.2004.07.003. MR2129092
- [8] P. W. Bates, J. Han, and G. Zhao, *On a nonlocal phase-field system*, Nonlinear Anal. **64** (2006), no. 10, 2251–2278, DOI 10.1016/j.na.2005.08.013. MR2213900
- [9] J. Cahn and J. Hilliard, *Free energy of a nonuniform system. I. Interfacial free energy*, J. Chem. Phys. **28** (1958), 258.
- [10] Q. Du, M. Gunzburger, R. B. Lehoucq, and K. Zhou, *Analysis and approximation of nonlocal diffusion problems with volume constraints*, SIAM Rev. **54** (2012), no. 4, 667–696, DOI 10.1137/110833294. MR3023366
- [11] Q. Du, L. Ju, X. Li, and Z. Qiao, *Stabilized linear semi-implicit schemes for the nonlocal Cahn-Hilliard equation*, J. Comput. Phys. **363** (2018), 39–54, DOI 10.1016/j.jcp.2018.02.023. MR3784415
- [12] Q. Du, L. Ju, X. Li, and Z. Qiao, *Maximum principle preserving exponential time differencing schemes for the nonlocal Allen-Cahn equation*, SIAM J. Numer. Anal. **57** (2019), no. 2, 875–898, DOI 10.1137/18M118236X. MR3945242
- [13] Q. Du and J. Yang, *Asymptotically compatible Fourier spectral approximations of nonlocal Allen-Cahn equations*, SIAM J. Numer. Anal. **54** (2016), no. 3, 1899–1919, DOI 10.1137/15M1039857. MR3514714
- [14] C. Duan, C. Liu, C. Wang, and X. Yue, *Convergence analysis of a numerical scheme for the porous medium equation by an energetic variational approach*, Numer. Math. Theory Methods Appl. **13** (2020), no. 1, 63–80, DOI 10.4208/nmtma.oa-2019-0073. MR4044415
- [15] W. E and J.-G. Liu, *Projection method. I. Convergence and numerical boundary layers*, SIAM J. Numer. Anal. **32** (1995), no. 4, 1017–1057, DOI 10.1137/0732047. MR1342281
- [16] W. E and J.-G. Liu, *Projection method. III. Spatial discretization on the staggered grid*, Math. Comp. **71** (2002), no. 237, 27–47, DOI 10.1090/S0025-5718-01-01313-8. MR1862987
- [17] D. J. Eyre, *Systems of Cahn-Hilliard equations*, SIAM J. Appl. Math. **53** (1993), no. 6, 1686–1712, DOI 10.1137/0153078. MR1247174
- [18] P. Fife, *Some Nonclassical Trends in Parabolic and Parabolic-like Evolutions*, Trends in Nonlinear Analysis, Springer, Berlin, 2003, pp. 153–191. MR1999098
- [19] H. Gajewski and K. Zacharias, *On a nonlocal phase separation model*, J. Math. Anal. Appl. **286** (2003), no. 1, 11–31, DOI 10.1016/S0022-247X(02)00425-0. MR2009615
- [20] C. G. Gal, A. Giorgini, and M. Grasselli, *The nonlocal Cahn-Hilliard equation with singular potential: well-posedness, regularity and strict separation property*, J. Differential Equations **263** (2017), no. 9, 5253–5297, DOI 10.1016/j.jde.2017.06.015. MR3688414
- [21] G. Giacomin and J. L. Lebowitz, *Phase segregation dynamics in particle systems with long range interactions. II. Interface motion*, SIAM J. Appl. Math. **58** (1998), no. 6, 1707–1729, DOI 10.1137/S0036139996313046. MR1638739
- [22] Z. Guan, J. Lowengrub, and C. Wang, *Convergence analysis for second-order accurate schemes for the periodic nonlocal Allen-Cahn and Cahn-Hilliard equations*, Math. Methods Appl. Sci. **40** (2017), no. 18, 6836–6863, DOI 10.1002/mma.4497. MR3742100
- [23] Z. Guan, J. S. Lowengrub, C. Wang, and S. M. Wise, *Second order convex splitting schemes for periodic nonlocal Cahn-Hilliard and Allen-Cahn equations*, J. Comput. Phys. **277** (2014), 48–71, DOI 10.1016/j.jcp.2014.08.001. MR3254224
- [24] Z. Guan, C. Wang, and S. M. Wise, *A convergent convex splitting scheme for the periodic nonlocal Cahn-Hilliard equation*, Numer. Math. **128** (2014), no. 2, 377–406, DOI 10.1007/s00211-014-0608-2. MR3259494
- [25] D. J. Horntrop, M. A. Katsoulakis, and D. G. Vlachos, *Spectral methods for mesoscopic models of pattern formation*, J. Comput. Phys. **173** (2001), no. 1, 364–390, DOI 10.1006/jcph.2001.6883. MR1857628

- [26] D. Li and Z. Qiao, *On second order semi-implicit Fourier spectral methods for 2D Cahn-Hilliard equations*, J. Sci. Comput. **70** (2017), no. 1, 301–341, DOI 10.1007/s10915-016-0251-4. MR3592143
- [27] D. Li and Z. Qiao, *On the stabilization size of semi-implicit Fourier-spectral methods for 3D Cahn-Hilliard equations*, Commun. Math. Sci. **15** (2017), no. 6, 1489–1506, DOI 10.4310/CMS.2017.v15.n6.a1. MR3668944
- [28] D. Li, Z. Qiao, and T. Tang, *Characterizing the stabilization size for semi-implicit Fourier-spectral method to phase field equations*, SIAM J. Numer. Anal. **54** (2016), no. 3, 1653–1681, DOI 10.1137/140993193. MR3507555
- [29] Z. Qiao and S. Sun, *Two-phase fluid simulation using a diffuse interface model with Peng-Robinson equation of state*, SIAM J. Sci. Comput. **36** (2014), no. 4, B708–B728, DOI 10.1137/130933745. MR3246906
- [30] R. Samelson, R. Temam, C. Wang, and S. Wang, *Surface pressure Poisson equation formulation of the primitive equations: numerical schemes*, SIAM J. Numer. Anal. **41** (2003), no. 3, 1163–1194, DOI 10.1137/S0036142901396284. MR2005199
- [31] R. Samelson, R. Temam, C. Wang, and S. Wang, *A fourth-order numerical method for the planetary geostrophic equations with inviscid geostrophic balance*, Numer. Math. **107** (2007), no. 4, 669–705, DOI 10.1007/s00211-007-0104-z. MR2342648
- [32] J. Shen, T. Tang, and L.-L. Wang, *Spectral Methods, Algorithms, Analysis and Applications*, Springer Series in Computational Mathematics, vol. 41, Springer, Heidelberg, 2011. MR2867779
- [33] J. Shen and X. Yang, *Numerical approximations of Allen-Cahn and Cahn-Hilliard equations*, Discrete Contin. Dyn. Syst. **28** (2010), no. 4, 1669–1691, DOI 10.3934/dcds.2010.28.1669. MR2679727
- [34] R. Temam, *Navier-Stokes Equations, Theory and Numerical Analysis*, AMS Chelsea Publishing, Providence, RI, 2001. Reprint of the 1984 edition. MR1846644
- [35] X. Tian and Q. Du, *Analysis and comparison of different approximations to nonlocal diffusion and linear peridynamic equations*, SIAM J. Numer. Anal. **51** (2013), no. 6, 3458–3482, DOI 10.1137/13091631X. MR3143839
- [36] X. Tian and Q. Du, *Asymptotically compatible schemes and applications to robust discretization of nonlocal models*, SIAM J. Numer. Anal. **52** (2014), no. 4, 1641–1665, DOI 10.1137/130942644. MR3231986
- [37] L. N. Trefethen, *Spectral methods in MATLAB*, Software, Environments, and Tools, vol. 10, Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 2000. MR1776072
- [38] C. Wang and J.-G. Liu, *Convergence of gauge method for incompressible flow*, Math. Comp. **69** (2000), no. 232, 1385–1407, DOI 10.1090/S0025-5718-00-01248-5. MR1710695
- [39] C. Wang and J.-G. Liu, *Analysis of finite difference schemes for unsteady Navier-Stokes equations in vorticity formulation*, Numer. Math. **91** (2002), no. 3, 543–576, DOI 10.1007/s002110100311. MR1907870
- [40] C. Wang, J.-G. Liu, and H. Johnston, *Analysis of a fourth order finite difference method for the incompressible Boussinesq equations*, Numer. Math. **97** (2004), no. 3, 555–594, DOI 10.1007/s00211-003-0508-3. MR2059469
- [41] L. Wang, W. Chen, and C. Wang, *An energy-conserving second order numerical scheme for nonlinear hyperbolic equation with an exponential nonlinear term*, J. Comput. Appl. Math. **280** (2015), 347–366, DOI 10.1016/j.cam.2014.11.043. MR3296065
- [42] S. M. Wise, C. Wang, and J. S. Lowengrub, *An energy-stable and convergent finite-difference scheme for the phase field crystal equation*, SIAM J. Numer. Anal. **47** (2009), no. 3, 2269–2288, DOI 10.1137/080738143. MR2519603
- [43] C. Xu and T. Tang, *Stability analysis of large time-stepping methods for epitaxial growth models*, SIAM J. Numer. Anal. **44** (2006), no. 4, 1759–1779, DOI 10.1137/050628143. MR2257126
- [44] K. Zhou and Q. Du, *Mathematical and numerical analysis of linear peridynamic models with nonlocal boundary conditions*, SIAM J. Numer. Anal. **48** (2010), no. 5, 1759–1780, DOI 10.1137/090781267. MR2733097

DEPARTMENT OF APPLIED MATHEMATICS, THE HONG KONG POLYTECHNIC UNIVERSITY, HUNG
HOM, KOWLOON, HONG KONG

Email address: `xiao11i@polyu.edu.hk`

DEPARTMENT OF APPLIED MATHEMATICS, THE HONG KONG POLYTECHNIC UNIVERSITY, HUNG
HOM, KOWLOON, HONG KONG

Email address: `zqiao@polyu.edu.hk`

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF MASSACHUSETTS, NORTH DARTMOUTH,
MASSACHUSETTS

Email address: `cwang1@umassd.edu`